



Version 2, July 2024

Authors: H.B. (Heike) Axmann MSc, dr. X. (Xuezhen) Guo, dr.ir. J. (Jan) Broeze, Charlotte Harbers MSc, dr. M. (Marcela) Viquez-Zamora, dr.ir. JM (Han) Soethoudt

Mitigate+: Food Loss and Waste country profile Colombia

Estimates of Food Loss and Waste, associated GHG emissions, nutritional losses, land use and water footprints

Urgency and call for action on FLW reduction

Theoretically, the world produces enough food to nourish the growing world population. Although precise data remains scarce, according to most recent studies, globally each year possibly as much as 30 per cent of the food produced is being lost or wasted somewhere between farm and fork. This not only represents a threat to food security but also severely and negatively impacts our food systems and natural resources. Food Loss and Waste (FLW) accounts for around 8 to 10 percent of our global Greenhouse Gas Emissions (GHGEs). Approximately a quarter of all freshwater used by agriculture is associated to the lost and wasted food. 4.4 million km² of land is used to grow food which is lost or wasted (FAO, 2019; WWF, 2021; Guo et al., 2020). The Sustainable Development Goal (SDG) Target 12.3 calls to 'halve per capita global food waste at the retail and consumer levels and reduce food losses along production and supply chains, including post-harvest losses' (Lipinski, B. 2022). With only 7 years to go, the world is far from being on track to achieve this target.

Way forward reducing FLW without baseline data

The UN and the Champions 12.3 Coalition launched the 'Target-Measure-Act approach' calling on all governments and companies to set FLW reduction targets, measure FLW, identify hotspots¹, and to take action to reduce FLW accordingly (Lipinski, 2020). However, with respect to primary data on FLW, much remains to be done. Just a

Food Loss and Waste (FLW) definition

FLW refers to all food intended for human consumption that is finally not consumed by humans. Food Loss is the decrease in the quantity or quality of food resulting from decisions and actions by food suppliers from the production stage in the chain, excluding retail, food service providers and consumers. Food Waste is the decrease in the quantity or quality of food resulting from decisions and actions by retailers, food services and consumers (FAO, 2019). Under this definition, FLW does not include food that is consumed in excess of nutritional requirements nor food that incurs a decrease of market value due to over-supply or other market forces, and not due to reduced quality.

1 In this document hotspots are defined as food products or food (sub) categories, eventually in combination with a supply chain link, that show the highest scores with respect to a selected (sub)set of sustainability indicators: FLW, GHGEs, nutrition, land use and water footprint.

2 Mutton & Goat Meat is not considered for the land-use footprints due to the marginal land consideration.

handful of mainly western countries have taken action to systematically measure and reduce FLW. Lack of data make it particularly difficult for lower-and-middle-income countries (LMIC), including Colombia, to specify the hotspot food products and chain stages, to define smart targets and to identify adequate interventions.

In order to contribute to this essential information we developed and used a mass flow model based on secondary data to derive the volume of FLW and the associated parameters accordingly (Guo et al., 2020). This approach allows to present an indicative country profile showing per food product category and chain stage not only the amount of FLW but also the GHGEs, the land-use and water footprints related to producing the FLW as well as induced nutrient losses. The sums differs per product and chain stage. Focusing on food products and chain stages which largely contribute to the aforementioned parameters can substantially lead to resource use efficiency and at the same time to climate mitigation action and nutrition security. This integrated approach towards FLW reduction can support policy makers and other food system actors taking informed decisions contributing to multiple sustainability objectives in parallel.

Modelling country data on FLW and FLW-associated GHGEs, land-use and water footprints and nutritional losses

FLW data was generated through a bottom-up, mass-flow model (Guo et al., 2020) that combines data on production and outputs as well as imports and exports at the country level. Estimates of losses per chain stage are derived from

Porter et al. (2016) to calculate the FLW in the supply chain according to the country's production and trade. The FLW-associated GHG emissions are calculated by using the GHG emission factors derived from Porter et al. (2016) to multiply the FLW at different supply chain stages.

Furthermore, a Protein and Nutrition Database developed by WUR (built on nutritional compositions derived from databases from FAO, USDA, Denmark and Japan) was used to calculate the nutritional value of the total consumed food in each country. The nutrient intakes are compared with estimated nutrition requirements per country (which is based on the composition of the population and per capita nutrient demand, according to WHO dietary recommendations).

In calculating the land use footprint of plant-based food items, FAO's 'Crops and livestock products' database is utilized by combining data on yields and harvested areas. This gives a simplified estimate of how much cropland is needed to grow the produce. Country-specific land use estimates for animal-based food items are however scarce. Therefore, global seaestimates as published by Poore & Nemecek (2018) are used. Applying this non-differentiated data has a drawback that it not accurately takes into account country-specific farming practices. Lastly, for the water footprint the broadly recognized datasets of Mekonnen and Hoekstra are used. These cover the Green, Blue and Grey water footprint of crops and derived crop products (Mekonnen & Hoekstra, 2011), and of animals and animal products (Mekonnen & Hoekstra, 2010).

FLW for Colombia- Top 15 Items




Figure 1 Top 15 hotspot categories of food loss and waste in terms of volumes and FLW-associated GHG emissions (in CO₂-eq.).



Land-use footprints of FLW for Colombia - Top 15 Items




Figure 2 Top 15 hotspot categories of the land-use footprints of FLW (in ha)

FLW, GHGs, nutrition, land use and water footprint country profile Colombia

Based on the country data modelling, estimates on FLW-associated GHGs were retrieved for Colombia and plotted with the FLW total tonnage to visualize the two components (Figure 1). For FLW the five main hotspots products are: fruits (others), bananas, milk, palm kernels, and plantains. However, ranking food categories according to the production of FLW-associated GHGs the five hotspot food products for Colombia are: bovine meat, milk, rice, poultry meat, and palm kernels. From the bovine meat chains, 0.13 million tons of FLW represents 4.7 million tons CO₂-eq. of GHGs. For milk, 3 million tons CO₂-eq. GHGs are generated from 1.2 million tons FLW. For rice 0.77 million FLW tons are generated, inducing 2 million tons CO₂-eq. GHGs.

Figure 2 presents the top 15 items with the largest land-use footprints of FLW. Animal-based products are dominating with bovine meat, milk, and poultry meat, ranking the top 3. Fruits, plantains and rice are the most land-use intensive plant-based individual items, ranking 4th and 5th and 6th.

With respect to the water footprints of the FLW, it is not dominated by either animal or plant-based items but demonstrate a mixed situation. Milk and plantains are the top 2 individual-item contributors followed by rice, bovine meat and poultry meat (Figure 3).

From another perspective, taking the percentages of FLW in relation to production percentages, vegetables and fruits are identified as the main hotspots showing average FLW 55% along the chains (Figure 4).



Water footprints of FLW for Colombia - Top 15 Items




Figure 3 Top 15 hotspot categories of the water footprints of FLW (in m3)

Colombia: % FLW/production for top 15 items




Figure 4 Percentages of FLW per product category

Further insights in hotspots are derived from estimated distribution of the FLW along supply chains in the number one hotspot product in the region (Figure 5). These data suggest that the agricultural production, processing & packaging, and postharvest handling & storage stages of the fruit and vegetable products are bottlenecks. These are focus points for more detailed data collection and analysis of causes to address potential interventions. Smart interventions in such 'hotspots' in food supply chains can

substantially contribute to GHG emission mitigation of food systems. Analysis of specificities of such chains (e.g. comparing informal and formal supply chains, and urban and rural settings) including comparison with supply chains for similar product categories may reveal promising interventions. Interventions may combine hardware (packaging, cooling, etc.), orgware (e.g. arrangements in chains) and software (knowledge, information) elements.

Colombia: FLW distribution (% of production) per chain stage for hotspot product categories




Figure 5 Percentages of FLW per stage in the supply chain for the number one hotspot product category.

Remark: Agricultural production does not include any potential yield gaps and focuses on actual production and harvest losses.



FLW-protein for Colombia - Top 15 Items

FLW associated Protein (1kg*10)




Figure 6 Top 15 hotspot categories of loss of proteins associated with FLW

Colombia - Nutrient supply (% of nutrient requirement)




Figure 7 Average provision of nutrients per capita relative to WHO dietary recommendations

Remark: because of uneven distribution of food over the population, parts of the population will suffer more insufficiencies than this diagram implies.

Figure 6 shows the protein losses associated with FLW where poultry meat, rice, milk, and bovine meat present again on the list of top 5 items. Finally, the food supply and FLW data were used to assess nutrient supply per capita in the Colombian population in relation to recommended nutrient intake (Figure 7). These are average numbers, and it is not likely that nutrients are evenly distributed across Colombia. Hence, there will be parts of the populations that suffer insufficiencies of calcium, folate, iron, vitamin A, and zinc. From nutrition security perspective, efforts for mitigating FLW in milk and rice chains would contribute the most to population nutrient gains (Table 1).

Table 1 Food product categories for which the FLW have highest share for the most critical nutrients.

Critical nutrients	FLW categories with highest loss of the nutrient (highest first)
Calcium	Milk, plantains, soybeans, rice, potato, onions, eggs, wheat
Folate	Rice, bananas
Iron	Rice, potato, wheat
Vitamin A	Poultry Meat, plantains, milk, eggs
Zinc	Rice, wheat, bovine meat, poultry, milk

Value loss

In Dep. Nac. Plan. (2016) the value loss for food waste (90 kton) in retail was identified at 204 billion pesos (64 million USD). The total value loss for Colombia was estimated at 5.4 billion USD².

Validation

The data used in Porter are not sufficiently granular to apply separate country commodity-stage-loss factors. Hence, the mass flow model described here might show data that don't match reality for the specific country at hand: Colombia. In that context a literature review was carried out to check whether the derived data are more or less in line with research related to Colombia itself. From the UN Food System Summit in 2021 it is known that national FLW data are scarce and difficult to obtain. Countries have been invited by the UN to take action on national level and develop policies laid down in a 'national pathway' document.

One article was found on FLW in Colombia as a whole. In a document from Dep. Nac. Plan. (2016) the FLW weight was identified for various categories, and can be compared to the data shown earlier, although the years they relate to are different.

Table 2 FLW weight (MTons) validation

Food category	FLW (here)	FLW (2016)
Fruits and vegetables	6.1	7.4
Roots and tubers	1	2.4
Cereals	1.3	0.77
Meat	0.55	0.27
Dairy	1.34	0.029

The table shows that cereals and dairy are overestimated, whereas for fruit and vegetables and roots and tubers the opposite holds. In general the top 3 FLW food categories are the same and more or less in line (although a factor 2 might be considered significant), and will lead to the correct food groups for FLW interventions.

According to Dep. Nac. Plan. (2016) 9.8 million tons (or 34%) of food destined for human consumption is lost and/or wasted. This is lower than the 13.5 million tons estimated from the mass-flow model that is used here.

Overall conclusions and suggestions for the next steps

Figure 8 displays a comprehensive ranking of hotspot food products based on five criteria. While there are nine hotspot food products identified, a closer examination reveals notable variations in the ranking of the nine hotspot products across different categories. Milk emerges as an extremely critical product, serving as a hotspot for all five categories. Bovine meat falls into the same category, positioned as a hotspot for four categories and represents the top hotspot product for GHGEs and land use footprints. In the next tier of hotspot products, rice and poultry meat both stand out among the top five hotspots for four categories and are classified as very critical hotspot products. Plantain is identified as a hotspot for three product categories and is classified as a critical product. Meanwhile, palm kernel holds the hotspot designation for two categories and is classified as moderately critical. Banana, despite being the most critical product in terms of FLW in tons, does not rank among the top five products in any of the other four categories and falls into the category of slightly critical products. Similarly, wheat and potatoes each hold a rank in only one hotspot category, both falling into the same category of slightly critical products, with wheat showing significant FLW-associated protein losses and potatoes for FLW tons.

It is suggested to develop FLW reduction actions, with synergy on GHGEs mitigation, nutrition, land-use and water footprints. The above analysis underlines that, if one considers sustainability in the context of these five selected indicators, the greatest impact can be achieved by concentrating efforts on milk, bovine meat, rice, and poultry compared to focusing on other food products.

Since the results are not on product level, it is not immediately clear, where to start your intervention. Our suggestion to develop FLW reduction actions, with synergy on GHGEs mitigation, nutrition, land-use and water footprints, is to implement monitoring or/and gather primary data for hotspot-supply chains of the country. The results in this document guide stakeholders by focusing on the top four food (sub) categories in combination with the indicative results on FLW per supply chain link. To research interventions, it is necessary to go to product level, which can be based on production or trade data in the country. The next step is to identify business cases for FLW reduction. For this purpose, WUR's EFFICIENT protocol³ and FLW cause and intervention tool⁴ can be used.

2 <https://www.foodandlandusecoalition.org/country/colombia/#:~:text=Approximately%20one%2Dthird%20of%20all,a%20healthy%20and%20active%20life.>

3 <https://edepot.wur.nl/556214> and <https://sites.google.com/iastate.edu/phlfrreduction/home/efficient-food-loss-waste-protocol>

4 The FLW cause & intervention tool (the-efficient-protocol.azurewebsites.net)

Colombia: Hotspot food products evaluated across five criteria




Figure 8 Ranking of hotspot product across five criteria

References

- Departamento Nacional de Planeación, Pérdida y desperdicio de alimentos en Colombia. 2016
- FAO. 2019. The State of Food and Agriculture 2019. Moving forward on food loss and waste reduction. Rome. Licence: CC BY-NC-SA 3.0 IGO.
- FAO, I., UNICEF, WFP and WHO., In Brief to The State of Food Security and Nutrition in the World 2021 – Transforming food systems for food security, improved nutrition and affordable healthy diets for all. 2021: p. 40.
- Guo, X.; Broeze, J.; Groot, J.J.; Axmann, H.; Vollebregt, M. 2020. A Worldwide Hotspot Analysis on Food Loss and Waste, Associated Greenhouse Gas Emissions, and Protein Losses. *Sustainability*, 12, 7488.
- Lipinski, B., Champions 12.3: SDG Target 12.3 on food loss and waste: 2022 progress report. 2022: p. 12.
- Mekonnen, M., & Hoekstra, A. Y. (2010). The green, blue and grey water footprint of animals and animal products. (Value of water research report 48; No. 48).
- Mekonnen, M. M., & Hoekstra, A. Y. (2011). The green, blue and grey water footprint of crops and derived crop products. *Hydrology and Earth System Sciences*, 15(5), 1577-1600.
- Poore, J., & Nemecek, T. (2018). Reducing food's environmental impacts through producers and consumers. *Science*, 360(6392), 987-992.
- Porter, S. D., Reay, D. S., Higgins, P., & Bomberg, E. 2016. A half-century of production-phase greenhouse gas emissions from food loss & waste in the global food supply chain. *Science of the Total Environment*, 571, 721-729.
- WWF, Driven to waste: The global impact of food loss and waste on farms. 2021: p. 24.

Colophon

Project number

623923390

Carried out by

The research that is documented in this study reports on work carried out by Wageningen Food & Biobased Research under Mitigate+ in 2022-2024. It was conducted in an objective way by the researchers.

Authors

Authors are researchers at Wageningen Food and Biobased Research, Wageningen, The Netherlands. Corresponding author: heike.axmann@wur.nl.

Acknowledgment

Support for this study was provided through Mitigate+ Initiative for Low-Emission Food Systems. We would like to thank all funders who supported this research through their contributions to the CGIAR Trust Fund.

Version

July 2024

